arXiv:nucl-ex/0405009v1 13 May 2004

ANISOTROPIC FLOW IN ULTRA-RELATIVISTIC HEAVY-ION COLLISIONS

MARKUS D. OLDENBURG
(for the STAR collaboration)
Ernest Orlando Lawrence Berkeley National Laboratory
One Cyclotron Road, Berkeley, CA 94720, USA
E-mail: MDOldenburg@lbl.gov

The analysis of anisotropic flow of particles created in high energy heavy-ion collisions gives insight into the early stage of these reactions. Measurements of directed flow (v_1), elliptic flow (v_2) and flow of 4^{th} and 6^{th} order (v_4 and v_6) are presented. While the study of v_2 for multi-strange particles establishes partonic collectivity the results for higher order anisotropies constrain the initial conditions of hydrodynamic model calculations.

1. Heavy-ion collisions

High-energy collisions of heavy ions provide a tool to study nuclear matter under extreme conditions. In the collision of two relativistic nuclei, a hot and dense system of partonic matter may form. During the formation time (of the order of $\tau_0 \sim 1\,\text{fm}/c$), partons from the colliding nuclei suffer hard scatterings. High-p_T hadrons are predominantly produced in these hard scatterings and, therefore, carry information about this early stage. If the system is thermalized and the relevant degrees of freedom are partonic then the state of matter created is referred to as the quark-gluon plasma. Due to a rapid expansion the system cools and at a critical temperature T_{c} undergoes a phase transition to hadronic matter[1]. This critical temperature T_{c} might be close to the temperature of chemical freeze-out T_{ch}, at which the last inelastic scatterings occur and particle yields no longer evolve. Thus hadron abundances provide information about the system at T_{ch} ($\sim T_{\text{c}}$). The system further expands and elastic scattering ceases. The corresponding temperature is called the kinetic freeze-out temperature T_{fo}. After kinetic freeze-out the momenta of the produced particles are fixed. Therefore the momentum distributions reflect the properties of the system at T_{fo}.

In summary one has to note that the discussed signals of the early and most interesting stage of the system's evolution are carried by only a tiny fraction of all produced particles, namely those with high transverse momentum p_T.

2. Anisotropic flow

In non-central heavy-ion collisions the overlap region of the two colliding nuclei is not azimuthally symmetric in the transverse plane. This leads to a larger pressure gradient in the reaction plane (defined by the incident beam direction and the impact parameter) than out-of-plane. These pressure gradients transform the original spatial anisotropy of the system (which is out-of-plane) in an in-plane anisotropy in momentum space. During this evolution the spatial anisotropy diminishes and the process of generating anisotropies in momentum space quenches itself. The measurement of the final azimuthal momentum anisotropy is therefore a signal which is carried by all particles, points back to the early evolution times of the collision, and provides information about the equation of state of the generated type of matter.

The transverse momentum asymmetry is analyzed utilizing a Fourier decomposition of the momentum distribution in each event with respect to the reaction plane[2]. By doing this one extracts Fourier coefficients of different order n: $v_n \equiv \langle \cos(n \cdot \phi) \rangle$, which characterize the strength of different orders of flow. v_1 and v_2 are called directed and elliptic flow, respectively.

3. The STAR experiment

The STAR experiment[3] at the Relativistic Heavy Ion Collider (RHIC) measures charged hadrons over a wide range in pseudorapidity η and transverse momentum p_T. Its main detector is a large TPC, sitting in a magnetic field of 0.5 T. Within the acceptance of $|\eta| < 1.0$ particles are identified by specific energy loss $\mathrm{d}E/\mathrm{d}x$. The two Forward-TPCs (FTPCs) add the region of $2.5 < |\eta| < 4.0$ to the overall acceptance. Other detector subsystems, which were not used in the studies shown here, include a silicon vertex tracker, an electromagnetic calorimeter and time-of-flight patches.

All results presented here were obtained using Au + Au collisions at a center of mass energy of $\sqrt{s_{NN}} = 200\,\mathrm{GeV}$.

4. Elliptic flow v_2

In addition to the large overall elliptic flow v_2 of several percent[4], which suggests a strongly anisotropic collective expansion, the measurement of v_2 shows a strong dependence on p_T (see Fig. 1, left). The almost linear rise of $v_2(p_T)$ up to $p_T \sim 1.5\,\mathrm{GeV}/c$ can for the first time be reproduced by hydrodynamic models[5], which overpredict elliptic flow at lower collision energies. These models assume local thermal equilibrium. The mass dependence of $v_2(p_T)$ at low p_T agrees with these models as well, with lighter particles experiencing more elliptic flow. This is consistent with all particles flowing within the same velocity field, since heavier particles have larger momentum for the same velocity.



Figure 1. Elliptic flow v_2 as a function of transverse momentum p_T. The left panel shows v_2 for particles with different strangeness content, together with model predictions. On the right the measurements are rescaled by the number of constituent quarks.

STAR measured the elliptic flow $v_2(p_T)$ of the multi-strange baryons $\Xi+\overline{\Xi}$ and $\Omega+\overline{\Omega}$ as well[6]. Notably, their anisotropic flow is comparable to the flow of the non-strange baryons. Since multi-strange baryons are expected to have small hadronic cross sections, it is unlikely that these particles could have picked up their large amount of flow during the hadronic stage only. Therefore this independence of strangeness content is a strong indication that collective motion is established during the partonic stage already.

At intermediate p_T ($1.5 < p_T < 5\,\mathrm{GeV}/c$) a clear ordering of mesons and baryons is visible. Quark coalescence models[7] predict that v_2 scaled by the number of constituent quarks n and plotted as a function of p_T/n will lie on top of each other. This universal curve for mesons and baryons represents v_2 developed by partons. As one can see from the measurement in Fig. 1 (right) quark coalescence in fact seems to be the dominant production mechanism at intermediate p_T.

5. Directed flow v_1 and the higher harmonics v_4 and v_6

The measurement of directed flow v_1 proved to be difficult due to its smallness at mid-rapidity. Therefore we utilized the Forward-TPCs[8] to determine v_1. As can be seen in Fig. 2 on the left panel significant directed flow is only visible in forward directions, while at mid-rapidity it is consistent with zero[9]. Comparisons to results on v_1 at lower beam energies look the same if plotted in the reference system of the corresponding beam rapidity.



Figure 2. Measurements of anisotropic flow of different orders. Left: Directed flow v_1 compared to results obtained by NA49. Right: Anisotropic flow of 4th and 6th order compared to the simple scaling of v_2 by $v_2^{n/2}$.

This measurement of directed flow v_1 was used to determine the sign of elliptic flow v_2 which was still undetermined but assumed to be positive[4]. The positiveness of v_2 could be confirmed[10] which is equivalent to the statement that elliptic flow is *in-plane*. In other words the simple description given in Sec. 2 holds and the initial spatial anisotropy indeed transforms into a anisotropy in momentum space which is rotated by 90°.

The strong elliptic flow made it possible to measure even higher orders of transverse anisotropy. While we measure sizable flow of 4th order, v_4, the signal of v_6 is consistent with zero (see Fig. 2, right)[9]. These results help to better constrain hydrodynamic model calculations, where the initial conditions manifest themselves predominantly in the higher orders of anisotropic flow[11]. Nevertheless, our measurements cannot be explained by these models so far. As another surprising fact it was noted that the higher harmonics of order n scale like $v_2^{n/2}$.

6. Conclusions

In summary we have shown that the measurement of anisotropic flow of different orders reveals interesting features of the early stage of the two colliding gold nuclei. Directed flow v_1 confirmed that elliptic flow v_2 is *in-plane* and first measurements of flow of higher orders give better constraints on hydrodynamic model calculations. While elliptic flow at low transverse momentum reproduces the mass scaling expected from hydrodynamic models, we observe a meson-baryon scaling at intermediate p_T. The flow of multi-strange baryons points to partonic collectivity.

Acknowledgments

We thank the RHIC Operations Group and RCF at BNL, and the NERSC Center at LBNL for their support. This work was supported in part by the HENP Divisions of the Office of Science of the U.S. DOE; the U.S. NSF; the BMBF of Germany; IN2P3, RA, RPL, and EMN of France; EPSRC of the United Kingdom; FAPESP of Brazil; the Russian Ministry of Science and Technology; the Ministry of Education and the NNSFC of China; Grant Agency of the Czech Republic, FOM and UU of the Netherlands, DAE, DST, and CSIR of the Government of India; the Swiss NSF.

References

1. F. Karsch, E. Laermann, A. Peikert, *Phys. Lett.* **B478**, 447 (2000).
2. S. A. Voloshin, Y. Zhang, *Z. Phys.* **C70**, 665 (1996).
3. K. H. Ackermann *et al.* (STAR collaboration), *Nucl. Instrum. Meth.* **A499**, 624 (2003).
4. K. H. Ackermann *et al.* (STAR collaboration), *Phys. Rev. Lett.* **86**, 402 (2001).
5. P. Huovinen *et al.*, *Phys. Lett.* **B503**, 58 (2001).
6. J. Castillo (for the STAR collaboration), to appear in the proceedings of Quark Matter 2004, *J. Phys.* **G**, *arXiv: nucl-ex/0403027* (2004).
7. D. Molnar, S. A. Voloshin, *Phys. Rev. Lett.* **91**, 092301 (2003); V. Grecco, C. M. Ko, P. Levai, *Phys. Rev.* **C68**, 034904 (2003); R. J. Fries, B. Müller, C. Nomaka, S. A. Bass, *Phys. Rev.* **C68**, 044902 (2003); Z. Lin, C. M. Ko, *Phys. Rev. Lett.* **89**, 202302 (2002).
8. K. H. Ackermann *et al.* (STAR collaboration), *Nucl. Instrum. Meth.* **A499**, 713 (2003).
9. J. Adams *et al.* (STAR collaboration), *Phys. Rev. Lett.* **92**, 062301 (2004).
10. M. D. Oldenburg (for the STAR collaboration), *arXiv: nucl-ex/0403007* (2004).
11. P. F. Kolb, *Phys. Rev.* **C68**, 031902(R) (2003).